

PRESSURE INJURY PREVENTION

Early & Targeted Intervention Kit (ETIK)



Pressure injuries are one of the largest unsolved healthcare challenges today 1-4

International pressure injury prevalence rates are estimated to be between:



IN LONG-TERM ACUTE CARE³

~27%

IN LONG-TERM CARE

3-32%

Why are pressure injuries difficult to prevent?



RAPID

Injuries can develop quickly with irreversible damage occurring in vulnerable patients within a matter of hours. 5,6



INVISIBLE

Inflammatory changes in the skin and underlying tissues may begin 3 to 10 days before becoming visible.⁷



VISUAL RISK ASSESSMENT TOOLS HAVE LIMITATIONS⁸

Visual risk assessment tools are based on subjective interpretation and have low predictive value. They do not always lead to effective pressure injury prevention.





Early & Targeted Intervention Kit (ETIK)

Supports you in aligning to the latest international guidelines.

EARLY

Ensure a 5-day*9 window of opportunity to deploy preventative intervention.

TARGETED

Targeted interventions guided by objective and anatomically-specific risk assessment.

INTERVENTION

Comprehensive and clinically proven interventions ranging from support surfaces, to repositioning and mobilisation.

KIT

Multi-component intervention kit, including clinical consultancy, supporting customer implementation.



Objective and early risk assessment with **Provizio® SEM Scanner**



Comprehensive **therapeutic support surfaces** portfolio



Targeted **repositioning** and enhanced **mobilisation**



Clinical consultancy delivering measurable and sustainable outcomes Elevate the pressure injury pathway in your care environment

 Ensure early and targeted risk assessment

 Deliver the appropriate intervention for your vulnerable patients

 Drive clinical excellence and best practice Scan the QR code or click the link below to get in touch with your Arjo Expert.

Visit www.arjo.com/ pressureinjury

Scan the QR code to view the product webpage for videos, specifications, training guide, gateway flyer and more.



www.arjo.com/ pressureinjury

Only Arjo designed parts, which are designed specifically for the purpose, should be used on the equipment and products supplied by Arjo. As our policy is one of continuous development we reserve the right to modify designs and specifications without prior notice. [®] and [™] are trademarks belonging to the Arjo group of companies. Copyright, trademarks and logos are the intellectual property of Bruin Biometrics LLC, this includes the following: Provizio[®] SEM Scanner and SEM Scanner

Bruin Biometrics LLCs copyright materials cannot be used or reproduced without Bruin Biometrics LLCs written consent. © Ario. 2021

At Arjo, we believe that empowering movement within healthcare environments is essential to quality care. Our products and solutions are designed to promote a safe and dignified experience through patient handling, medical beds, personal hygiene, disinfection, diagnostics, and the prevention of pressure injuries and venous thromboembolism. With over 6000 people worldwide and 60 years caring for patients and healthcare professionals, we are committed to driving healthier outcomes for people facing mobility challenges.

Arjo AB · Hans Michelsensgatan 10 · 211 20 Malmö · Sweden · +46 10 335 4500

www.arjo.com

* Median

References

1. Li Z, Lin F, Thalib L, Chaboyer W. Global prevalence and incidence of pressure injuries in hospitalised adult patients: A systematic review and meta-analysis. Int J Nurs Stud. 2020 May. 2. Moore Z, Avsar P, Conaty L, Moore D.H, Patton D, & O'Connor T (2019) The prevalence of pressure ulcers in Europe, what does the Europen data tell us? Journal of Wound Care. 3. Berlowitz D, Lukas CV, Parker V, Niederhauser A, Silver J, Logan Cet al. Preventing pressure ulcers in hospitals: a toolkit for improving quality of care [Internet]. Rockville (MD): Agency for Healthcare Research and Quality; 2014 [cited 2018 Oct 2]. 4. Anthony, D. M., Alosoumi, D., and Safari, R. (2019). Prevalence of pressure ulcers in long term care: A global review, Journal of Wound Care, 28(11), pp. 1-7. DOI: 10.12968/jowc.2019.28.11.702. 5. Gefen A (2018) The future of pressure ulcer prevention is here: Detecting and targeting inflammation early. EWMA Journal 2018, 19(2):7-13. 6. Gefen, A (2008). How much time does it take to get a pressure ulcer? Integrated evidence from human, animal and invitro studies. Ostomy Wound Manage. 2008b; 54(10): 26-8,30-5. 7. Moore Z, Patton D, Rhodes SL, O'Connor T. Subepidermal moisture and bioimpedance: a literature review of a novel method for early detection of pressure-induced tissue damage (pressure ulcers). Int Wound J 2017; 14:331-337. 8. Moore ZEH, Patton D. Risk assessment tools for the prevention of pressure ulcers. Cochrane Database of Systematic Reviews 2019, Issue 1. Art No.:CD006471.DOI:10.1002/14651858.CD006471.Pub.4. 9. Coknokwo H, Bryant R, Milne J et al. A blinded clinical study using subepidermal moisture biocapacitance measurement device for early detection of pressure injuries. Wound Repair & Reg 2020;1-11.

